



Photosynthetic Pigments of oceanic Chlorophyta belonging to prasinophytes clade VII

Journal:	<i>Journal of Phycology</i>
Manuscript ID:	Draft
Manuscript Type:	Note
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Lopes dos Santos, Adriana; Sorbonne Universités, UPMC Univ Paris 06, UMR 7144, Station Biologique, Place Georges Teissier; CNRS, UMR 7144, Station Biologique, Place Georges Teissier</p> <p>Gourvil, Priscillia; Sorbonne Universités, UPMC Univ Paris 06, UMR 7144, Station Biologique, Place Georges Teissier; CNRS, UMR 7144, Station Biologique, Place Georges Teissier</p> <p>Rodriguez, Francisco; Instituto Español de Oceanografía, Centro Oceanográfico de Vigo, Subida a Radio Faro</p> <p>Garrido, José; Instituto de Investigaciones Marinas(CSIC), Av. Eduardo Cabello, 6</p> <p>Vaulot, Daniel; Sorbonne Universités, UPMC Univ Paris 06, UMR 7144, Station Biologique, Place Georges Teissier; CNRS, UMR 7144, Station Biologique, Place Georges Teissier</p>
Keywords:	phytoplankton, picoplankton, prasinophytes, pigments, HPLC

Article Type: Research Note

Photosynthetic Pigments of oceanic Chlorophyta belonging to prasinophytes clade VII.

Adriana Lopes dos Santos^{1,2}, Priscillia Gourvil^{1,2}, Francisco Rodriguez-Hernandez³, José Luis Garrido⁴, and Daniel Vaultot^{1,2*}.

¹Sorbonne Universités, UPMC Univ. Paris 06, UMR 7144, Station Biologique, Place Georges Teissier, 29680 Roscoff, France

²CNRS, UMR 7144, Station Biologique, Place Georges Teissier, 29680 Roscoff, France

³Microalgas Nocivas (IEO), Unidad Asociada al IIM (CSIC). Instituto Español de Oceanografía (IEO), Centro Oceanográfico de Vigo, Subida a Radio Faro 36390 Vigo, España.

⁴Microalgas Nocivas (IEO), Unidad Asociada al IIM (CSIC). Instituto de Investigaciones Marinas(CSIC). Av. Eduardo Cabello, 6. 36208 Vigo, España.

Submitted to Journal of Phycology, April 25, 2015

* Author for correspondence : vaultot@gmail.com

Abstract

The ecological importance and diversity of pico/nano-planktonic algae remains poorly studied in marine waters, in part because many are tiny and without distinctive morphological features. Amongst green algae, Mamiellophyceae such as *Micromonas* or *Bathycoccus* are dominant in coastal waters while prasinophytes clade VII, yet not formerly described, appear to be major players in open oceanic waters. The pigment composition of 13 strains representative of different sub-clades of clade VII was analyzed using a method that improves the separation of loroxanthin and neoxanthin. All the prasinophytes clade VII analyzed here showed a pigment composition similar to that previously reported for RCC287 corresponding to pigment group prasino-2A. However we detected in addition astaxanthin for which it is the first report in prasinophytes. Among the strains analyzed the pigment signature are qualitatively similar within sub-clades A and B. In contrast RCC3402 from sub-clade C (*Picocystis*) lacks loroxanthin, astaxanthin and antheraxanthin. For sub-clades A and B, loroxanthin was lowest at highest light irradiance suggesting a light-harvesting role of this pigment in clade VII as in *Tetraselmis*.

Keywords : phytoplankton, picoplankton, prasinophytes, pigments, HPLC

The paraphyletic group of prasinophytes is an assemblage of free-living unicellular microalgae present in both marine and freshwater habitats (Leliaert et al. 2012). Molecular phylogenetic, ultra-structural, and biochemical approaches have helped taxonomists to reorganize gradually the group into new classes and clades (Guillou et al. 2004, Marin and Melkonian 2010, Subirana et al. 2013, Lemieux et al. 2014a). Currently the prasinophytes are divided into nine groups known as clades I to IX, based on phylogenetic analyses of the nuclear 18S (nuclear-encoded small subunit rRNA) gene (Fawley et al. 2000, Guillou et al. 2004, Viprey et al. 2008). These clades may correspond to true classes, or be composed of a small number of species or of environmental sequences only. For example, Chlorodendrophyceae (Massjuk 2006) known previously as prasinophytes clade IV was recently raised to the class level and added to the “core of chlorophytes” (Fucikova et al. 2014). Clade V corresponds to the order Pycnococaceae with two major species, *Pseudoscurfieldia marina* and *Pycnococcus provasolii* which are probably two forms of a single life cycle (Fawley et al. 1999, Guillou et al. 2004). Clades VIII and IX are composed entirely by environmental sequences without representatives in culture (Viprey et al. 2008). Clade II, previously corresponding to the order Mamiellales, was raised recently to the class level as Mamiellophyceae (Marin and Melkonian 2010) and contains three important genera of marine pico-phytoplankton: *Micromonas* (Butcher 1952), *Bathycoccus* (Eikrem and Throndsen 1990) and *Ostreococcus* (Chrétiennot-Dinet et al. 1995).

In coastal waters, Mamiellophyceae appear largely dominant, especially within the pico-plankton, with the genus *Micromonas* making the highest contribution and followed to a lesser extent by *Bathycoccus* (Throndsen, J. and Kristiansen 1991, Not et al. 2004, Collado-Fabri et al. 2011, Balzano et al. 2012). In contrast in the open ocean, another group of prasinophytes, clade VII, with cell size in the 3 to 5 μm range, has been found to make an important contribution to

the pico-plankton community in regions such as the Equatorial Pacific and Mediterranean Sea (Moon-van der Staay et al. 2000, Viprey et al. 2008, Shi et al. 2009). The distribution of clade VII in typically oceanic mesotrophic waters makes this an interesting group. Prasinophyte clade VII contains several cultured strains, but it has not been described formerly yet. Guillou *et al.* (2004) divided this group into three well-supported sub-clades, A, B and C, the latter being formed by *Picocystis salinarum*, a small species found in saline lakes (Lewin et al. 2000, Roesler et al. 2002, Krienitz et al. 2012).

Traditionally, pigment signature has been used to determine the taxonomy of algae groups present in the water column (Jeffrey et al. 1997). This approach has been largely superseded by molecular approaches (Liu et al. 2009) but pigments remain an important phenotypic characteristic that allowed to point out the importance of green algae in specific regions of Pacific Ocean, Mediterranean Sea or Arctic Ocean (Obayashi and Tanoue 2002, Miki et al. 2008, Gutiérrez-Rodríguez et al. 2010, Coupel et al. 2014). The use of pigments in order to discriminate different types of prasinophytes has proven to be a difficult task since a diversity of photosynthetic signatures can be found in this group. Prasinophytes can be divided into three major groups based on their carotenoid composition (Egeland et al. 1997, Garrido et al. 2009). Group 1 contains the basic set of carotenoids present in Chlorophyceae: neoxanthin, violaxanthin, lutein, zeaxanthin, antheraxanthin, astaxanthin and β - β -carotene. The two later pigments are found to accumulate in *Haematococcus* and *Dunaliella* species under stress conditions (Lemoine and Schoefs 2010). Group 2 consists of the basic set of carotenoids plus loroxanthin (2A) and siphonaxanthin (2B). Group 3 contains prasinoxanthin (3A) and uriolide/micromonal (3B) in addition to the pigments found in group 2 (Jeffrey et al. 2011)

Within clade VII, a single strain RCC287 has been analyzed for its pigment composition (Latasa et al. 2004). A large number of clade VII strains are available from the Roscoff Culture Collection (<http://roscoff-culture-collection.org/>) originating from a range of environment. Their genetic diversity is currently analyzed allowing further delineation of new sub-clades (Lopes et al. in preparation) within the two major sub-clades A and B described by Guillou *et al* (2004). Among the 44 strains of clade VII present in the RCC, 39 belong to sub – clade A. The aim of this study was to determine the phenotypic characteristics of this important group of marine green algae by analyzing the pigment composition of thirteen strains belonging to the three sub-clades (A, B, C) of prasinophytes clade VII and isolated from a range of oceanic location and depths (Table 1). We also assessed the effect of three light irradiances on pigment composition for a subset of these strains.

Thirteen strains belonging to clade VII (Table 1) were grown at 22°C in K seawater medium (Keller et al. 1987) under 140 $\mu\text{E.m}^{-2}.\text{s}^{-1}$ in continuous light. A subset of nine strains was also grown at two other light levels (14 and 65 $\mu\text{E.m}^{-2}.\text{s}^{-1}$). Prior to sample collection, cell concentration was determined by flow cytometry using a Becton Dickinson Accuri C6. Approximately 50 ml of cultures were harvested in late exponential or early stationary phase and were filtered onto glass fiber GF/F (Whatman, Maidstone, UK) without vacuum and immediately frozen in liquid nitrogen and stored at -80°C. Pigments were analyzed within one month. Frozen filters were extracted with 3 mL of 90% acetone in screw cap glass tubes with polytetra-fluoroethylene (PTFE) lined caps, placed in an ice-water bath. After 15 minutes, filters were homogenized using a stainless steel spatula for filter grinding. Tubes were placed in an ultrasonic bath with water and ice for 5 minutes. The slurries were then centrifuged 5 minutes at 4.500 r.p.m. and supernatants filtered through 13 mm diameter polypropylene syringe filters (MS

PTFE, 0.22 μm pore size) to remove cell and filter debris. Before injection 1ml of each sample extract was added with 0.4 ml of Milli-Q water to avoid peak distortion. Pigments extracted from clade VII strains were analyzed using a modification of Zapata *et al.* (2000) method, described by Garrido *et al.* (2009) to improve the separation of loroxanthin and neoxanthin (Table S1). All graphs and analyses were performed with the R software using the ggplot2 and FactoMineR libraries (R Development Core Team 2013).

Intracellular chlorophyll (Chl) *a* content ranged from 4 to 26 fg per cell in most strains except for RCC996 (VIIA) and RCC3402 (*Picocystis* - clade VIIC) for which it was much higher (Table 2). This range agreed with values previously determined for marine microalgae in the same size range (Simon et al. 1994).

All the prasinophytes clade VIIA and B analyzed here showed a very similar pigment composition (Table 2). It did not seem to change drastically with between sub-clades A and B, nor with the depth of isolation (Figure 1). This composition is very similar to that reported for RCC287 by Latasa *et al.* (2004) corresponding to pigment group prasino-2A. We did not observe strong differences (Figure 1, Table 2) with the data of Latasa *et al.* (2004): in particular the ratios obtained for zeaxanthin and lutein, were very similar in both studies despite the slight difference in light levels (100 vs. 140 $\mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ in our study): zeaxanthin, 0.042 (w/w) *versus* 0.043 (w/w) and lutein, 0.382 (w/w) *versus* 0.363 (w/w). However their study used a less resolutive method and did not report the presence of loroxanthin and astaxanthin in RCC287. For loroxanthin this is probably due to the co-elution of this pigment with neoxanthin in the analytic method employed by these authors. For astaxanthin, our strains were grown in continuous light in contrast to Latasa *et al.* (2004) who used L:D cycles. This might have triggered the synthesis and accumulation of

this carotenoid, which is known to be synthesized under stressful conditions, such as high light, UV irradiation or nutrient depletion in Chlorophyceae, like *Haematococcus* and *Dunaliella* (Lemoine and Schoefs 2010) as well in some strains of Trebouxiophyceae such as *Picochlorum* (Lubián et al. 2000). In *Haematococcus*, maximal production of astaxanthin has been obtained under to continuous illumination (Domínguez-Bocanegra et al. 2004)

In our study, only RCC1124 (sub-clade A) did not contain loroxanthin within strains belonging sub-clades A and B (Table 2). *Picocystis* (RCC3402, clade VIIC) did not contain loroxanthin, astaxanthin and antheraxanthin (Table 1). Violaxanthin and lutein were the most abundant carotenoids for clade VIIA and VIIB. Astaxanthin came as the third more abundant within VIIA and RCC2337 (VIIB), while β - ϵ -carotene came third for RCC2339 (VIIB). For *Picocystis* (clade VIIC), lutein and β - ϵ -carotene were the most abundant carotenoids (Figure 1, Table 2), but the ratio of accessory pigments to Chl *a* was much lower than in clades VIIA and B (Figure 1).

We analyzed the influence of irradiance (14, 65 and 140 $\mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) on pigment composition of nine strains of prasinophytes VIIA and B (Figure 2 and S1, and Table S2). Change in light intensities induce opposite trends in the ratios of photosynthetic and photoprotective pigments relative to Chl *a*. Accessory chlorophylls and carotenoids involved in light capture either increase relative to Chl *a* at low light, while photoprotective carotenoids increase at high light (Schlüter et al. 2000, Henriksen et al. 2002). In our study, Chl *b* ratios increased slightly at low light, as expected, except for RCC3376 that showed a very slightly lower ratio at low light than at high light (0.78 vs. 0.81) (Figure 2 and Table S2). A similar slight decrease was also observed by Garrido *et al.* (2009) for the green alga *Tetraselmis suecia*.

Neoxanthin, β - β carotene and loroxanthin appeared to have a light harvesting role for most of the strains, except RCC287 (Figure 2 and Table S2), but change could be subtle as in the case of neoxanthin. Neoxanthin has been found to be associated with light harvesting complexes in the Mamiellophyceae *Mantoniella squamata* (Wilhelm and Lenarz-Weiler 1987). The most drastic changes were observed for loroxanthin (Figure 2) suggesting that this pigment has a major light harvesting role in clades VII A and B in agreement with what observed by Garrido *et al.* (2009) with *T. suecica*. One strain, RCC3376 (sub-clade A), had very low ratios of loroxanthin to Chl *a* which did not seem to change with light (Figure 2).

The increase of astaxanthin (from 2 to 4-fold depending on the strains) with light intensity suggests that this carotenoid has a photoprotective role for in clade VIIA and B (Figure 2), as previously demonstrated in the Chlorophyceae *Haematococcus pluvialis* (Wang et al. 2003, Gao et al. 2012). Among all strains, RCC3374 showed the most impressive accumulation of astaxanthin which contributed up to 42% of the total carotenoid pool under high light conditions (Figure 2). In comparison, *H. pluvialis* can accumulate 86 - 90% of astaxanthin in the total carotenoid pool after sixteen days cultures of under stress conditions (Sarada et al. 2002). However RCC3374 cell size (3 μ m) is about 7 times smaller than *H. pluvialis* resting cells (25 μ m, Gu et al. 2013) and maximum cell density is $3 \cdot 10^6$ cell mL⁻¹. A rough computation show that our RCC3374 culture was able to produce only 0.006 mg/L of astaxanthin against 8.3–10.69 mg/l found in old cultures of *H. pluvialis* (Sarada et al. 2002). If one could increase the cell concentration about 1000-fold with a different medium and manipulate astaxanthin cell content through stress, then clade VII could be an interesting contender for astaxanthin production.

Lutein also showed a photoprotective behavior. Its contribution to total carotenoids increased sharply from low to medium light and the stabilized at the highest light (Figure 2, Table S2). Such increase under high light conditions has been previously reported by Böhme et al. (2002) in the Mamiellophyceae *M. squamata*. These authors suggested that lutein played an important role as intermediate of biosynthesis for light harvesting pigments after light shifts from HL to LL. This role was coherent with its loose binding to the LHC apoprotein, also observed for the violaxanthin cycle (VAZ) carotenoids. However, lutein and loroxanthin are xanthophylls derived from β - ϵ carotene, and both have also been suggested also to take part in photoprotective mechanisms (non-photochemical quenching, NPQ) to prevent photo-oxidative damage in high light conditions in the green alga *Chlamydomonas reinhardtii* (Niyogi et al. 1997).

As for lutein, the content of the photoprotective xanthophyll cycle involving violaxanthin, antheraxanthin and zeaxanthin (VAZ cycle) relative to Chl *a* increased in general sharply from low to medium light and the stabilized at high light (Figure 2 and S1, Table S2). Their specific behavior however differed among strains. For example, zeaxanthin did not change much in RCC287 and RCC857 while it increased several-fold in other strains (e.g. RCC719, Figure 2).

In order to assess the visualize any hidden relationship between strain origin and pigment composition, we performed a Principal Component Analysis (PCA) based on pigments to Chl *a* ratios (Figure 3). The first two components explained more than 50% (dimension 1 and 2, 33.1 % and 20.4 %, respectively) of the variance. Pigments contributing positively to dimension 1 corresponded to photoprotective pigments (lutein, β - ϵ carotene, zeaxanthin, violaxanthin, antheraxanthin and astaxanthin) while pigments involved in photosynthetic capture, such as loroxanthin and β - β -carotene, contributed negatively to this axis. Pigments with moderate

response to light, such as Chl *b* and neoxanthin, contributed to dimension 2. Strains under medium and high light were separated from those at low light along dimension 1 which reflects that for several pigments the largest changes occurred between low and medium light and then stabilized at high light. This analysis confirmed that pigment composition and change did not seem to be linked to strain sub-clade (A vs. B) or depth of isolation (surface vs. DCM).

Recent phylogenetic results pointed clade VII A/B as a sister group of the core of Chlorophyta (Guillou et al. 2004, Leliaert et al. 2012, Lemieux et al. 2014a, 2014b). The presence of astaxanthin as in core Chlorophyta while it is absent with other prasinophytes may reflects another common feature between clade VII and core Chlorophyta. Moreover while Guillou *et al* (2004) included *Picocystis* into clade VII based on the phylogenetic analysis of 18S rRNA gene, the recent analysis of chloroplast genomes (Lemieux et al. 2014a) has shown widely divergent traits between *Picocystis* on one side and sub-clades VIIA and B, on the other side. The divergent carotenoid composition of *Picocystis* (absence of loroxanthin, astaxanthin, and antheraxanthin) reinforce these phylogenetic analyses.

Some of the important findings of this study include the discovery of loroxanthin and astaxanthin in prasinophytes clades VIIA and B, which had not been reported before, as well the possible role of the former pigment as a light harvesting accessory pigment. This should prompt the need to reexamine the pigment composition of other members of this diverse and ancient group constituted by prasinophytes using improved analytical protocols.

Acknowledgments

Financial support for this work was provided by the European Union program MaCuMBA (FP7-KBBE-2012-6-311975).

References

- Balzano, S., Marie, D., Gourvil, P. & Vaultot, D. 2012. Composition of the summer photosynthetic pico and nanoplankton communities in the Beaufort Sea assessed by T-RFLP and sequences of the 18S rRNA gene from flow cytometry sorted samples. *ISME J.* 6:1480–98.
- Böhme, K., Wilhelm, C. & Goss, R. 2002. Light regulation of carotenoid biosynthesis in the prasinophycean alga *Mantoniella squamata*. *Photochem. Photobiol. Sci.* 1:619–28.
- Butcher, R. 1952. Contributions to our knowledge of the smaller marine algae. *J. Mar. Biol. Assoc. U.K.* 19:175–91.
- Chrétiennot-Dinet, M.-J., Courties, C., Vaquer, A., Neveux, J., Claustre, H., Lautier, J. & Machado, M.C. 1995. A new marine picoeucaryote: *Ostreococcus tauri* gen. et sp. nov. (Chlorophyta, Prasinophyceae). *Phycologia*. 34:285–92.
- Collado-Fabri, S., Vaultot, D. & Ulloa, O. 2011. Structure and seasonal dynamics of the eukaryotic picophytoplankton community in a wind-driven coastal upwelling ecosystem. *Limnol. Oceanogr.* 56:2334–46.
- Coupe, P., Matsuoka, A., Ruiz-Pino, D., Gosselin, M., Claustre, H., Marie, D., Tremblay, J.E. et al. 2014. Pigment signatures of phytoplankton communities in the Beaufort Sea. *Biogeosciences Discuss.* 11:14489–530.
- Domínguez-Bocanegra, A.R., Guerrero Legarreta, I., Martínez Jerónimo, F. & Tomasini Campocósio, A. 2004. Influence of environmental and nutritional factors in the production of astaxanthin from *Haematococcus pluvialis*. *Bioresour. Technol.* 92:209–14.
- Egeland, E.S., Guillard, R.R.L. & Liaaen-Jensen, S. 1997. Additional carotenoid prototype representatives and a general chemosystematic evaluation of carotenoids in Prasinophyceae (Chlorophyta). *Phytochemistry*. 44:1087–97.
- Eikrem, W. & Throndsen, J. 1990. The ultrastructure of *Bathycoccus* gen. nov. and *B. prasinos* sp. nov., a non-motile picoplanktonic alga (Chlorophyta, Prasinophyceae) from the Mediterranean and Atlantic. *Phycologia*. 29:344–50.
- Fawley, M.W., Qin, M. & Yun, Y. 1999. The relationship between *Pseudoscurfieldia marina* and *Pycnococcus provasolii* (Prasinophyceae, Chlorophyta): Evidence from 18S rDNA sequence data. *J. Phycol.* 35:838–43.
- Fawley, M.W., Yun, Y. & Qin, M. 2000. Phylogenetic analyses of 18S rDNA sequences reveal a new coccoid lineage of the Prasinophyceae (Chlorophyta). *J. Phycol.* 36:387–93.
- Fucikova, K., Leliaert, F., Cooper, E.D., Åkaloud, P., D'Hondt, S., De Clerck, O., Gurgel, C.F.D. et al. 2014. New phylogenetic hypotheses for the core Chlorophyta based on chloroplast sequence data. *Front. Ecol. Evol.* 2.
- Gao, Z., Meng, C., Zhang, X., Xu, D., Zhao, Y., Wang, Y., Lv, H. et al. 2012. Differential expression of carotenogenic genes, associated changes on astaxanthin production and photosynthesis features induced by JA in *H. pluvialis*. *PLoS One*. 7:e42243.
- Garrido, J.L., Rodríguez, F. & Zapata, M. 2009. Occurrence of loroxanthin, loroxanthin decenoate, and loroxanthin dodecenoate in *Tetraselmis* species (Prasinophyceae, Chlorophyta). *J. Phycol.* 45:366–74.

252 Gu, W., Xie, X., Gao, S., Zhou, W., Pan, G. & Wang, G. 2013. Comparison of different cells of *Haematococcus*
253 *pluvialis* reveals an extensive acclimation mechanism during its aging process: from a perspective of
254 photosynthesis. *PLoS One*. 8:e67028.

255 Guillou, L., Eikrem, W., Chrétiennot-Dinet, M.-J., Le Gall, F., Massana, R., Romari, K., Pedrós-Alió, C. et al. 2004.
256 Diversity of picoplanktonic prasinophytes assessed by direct nuclear SSU rDNA sequencing of environmental
257 samples and novel isolates retrieved from oceanic and coastal marine ecosystems. *Protist*. 155:193–214.

258 Gutiérrez-Rodríguez, A., Latasa, M., Estrada, M., Vidal, M. & Marrasé, C. 2010. Carbon fluxes through major
259 phytoplankton groups during the spring bloom and post-bloom in the Northwestern Mediterranean Sea. *Deep.*
260 *Res. Part I Oceanogr. Res. Pap.* 57:486–500.

261 Henriksen, P., Riemann, B., Kaas, H., Sorensen, H.L.H.M. & Sørensen, H.M.H.L. 2002. Effects of nutrient-
262 limitation and irradiance on marine phytoplankton pigments. *J. Plankton Res.* 24:835–58.

263 Jeffrey, S., Wright, S. & Zapata, M. 2011. Phytoplankton Pigments: Characterization, Chemotaxonomy and
264 Applications in Oceanography. In Roy S, Egeland ES, Johnsen G, L. C. [Ed.]. Cambridge University Press,
265 Cambridge., pp. 3–77.

266 Jeffrey, S.W., Mantoura, R.F.C. & Wright, S. 1997. Phytoplankton Pigments in Oceanography: Guidelines to
267 Modern Methods. Unesco Publishing. 661 pp.

268 Keller, M.D., Selvin, R.C., Claus, W. & Guillard, R.R.L. 1987. Media for the culture of oceanic ultraphytoplankton.
269 *J. Phycol.* 23:633–8.

270 Krienitz, L., Bock, C., Kotut, K. & Luo, W. 2012. *Picocystis salinarum* (Chlorophyta) in saline lakes and hot springs
271 of East Africa. *Phycologia*. 51:22–32.

272 Latasa, M., Scharek, R., Le Gall, F. & Guillou, L. 2004. Pigment suites and taxonomic groups in Prasinophyceae. *J.*
273 *Phycol.* 40:1149–55.

274 Leliaert, F., Smith, D.R., Moreau, H., Herron, M.D., Verbruggen, H., Delwiche, C.F. & De Clerck, O. 2012.
275 Phylogeny and molecular evolution of the green algae. *CRC. Crit. Rev. Plant Sci.* 31:1–46.

276 Lemieux, C., Otis, C. & Turmel, M. 2014a. Six newly sequenced chloroplast genomes from prasinophyte green algae
277 provide insights into the relationships among prasinophyte lineages and the diversity of streamlined genome
278 architecture in picoplanktonic species. *BMC Genomics*. 15:857.

279 Lemieux, C., Otis, C. & Turmel, M. 2014b. Chloroplast phylogenomic analysis resolves deep-level relationships
280 within the green algal class Trebouxiophyceae. *BMC Evol. Biol.* 14:211.

281 Lemoine, Y. & Schoefs, B. 2010. Secondary ketocarotenoid astaxanthin biosynthesis in algae: a multifunctional
282 response to stress. *Photosynth. Res.* 106:155–77.

283 Lewin, R.A., Krienitz, L., Oericki, R.G., Takeda, H. & Hepperle, D. 2000. *Picocystis salinarum* gen. et sp. nov.
284 (Chlorophyta) - a new picoplanktonic green alga. *Phycologia*. 39:560–5.

285 Liu, H., Probert, I., Uitz, J., Claustre, H., Aris-Brosou, S., Frada, M., Not, F. et al. 2009. Haptophyta rule the waves:
286 extreme oceanic biodiversity in non-calcifying haptophytes explains the 19-Hex paradox. *Proc. Natl. Acad.*
287 *Sci.* 106:12803–8.

Lopes et al. Pigments in prasinophytes clade VII - 14

- 288 Lubián, L., Montero, O., Moreno-Garrido, I., Huertas, I.E., Sobrino, C., González-del Valle, M. & Pares, G. 2000.
289 *Nannochloropsis* (Eustigmatophyceae) as source of commercially valuable pigments. *J. Appl. Phycol.* 12:249–
290 55.
- 291 Marin, B. & Melkonian, M. 2010. Molecular phylogeny and classification of the Mamiellophyceae class. nov.
292 (Chlorophyta) based on sequence comparisons of the nuclear- and plastid-encoded rRNA Operons. *Protist.*
293 161:304–36.
- 294 Massjuk, N.P. 2006. Chlorodendrophyceae class. nov. (Chlorophyta, Viridiplantae) in the Ukrainian flora: I. The
295 volume, phylogenetic relations and taxonomical status. *Ukr. Bot. J.* 63:601–6014.
- 296 Miki, M., Ramaiah, N., Takeda, S. & Furuya, K. 2008. Phytoplankton dynamics associated with the monsoon in the
297 Sulu Sea as revealed by pigment signature. *J. Oceanogr.* 64:663–73.
- 298 Moon-van der Staay, S., Van der Staay, G.W.M., Guillou, L., Vaultot, D., Claustre, H. & Medlin, L.. 2000.
299 Abundance and diversity of prymnesiophytes in the picoplankton community from the equatorial Pacific
300 Ocean inferred from 18S rDNA sequences. *Limnol. Oceanogr.* 45:98–109.
- 301 Niyogi, K.K., Björkman, O. & Grossman, A.R. 1997. The roles of specific xanthophylls in photoprotection. *Proc.*
302 *Natl. Acad. Sci. U. S. A.* 94:14162–7.
- 303 Not, F., Latasa, M., Marie, D., Cariou, T., Vaultot, D. & Simon, N. 2004. A single species *Micromonas pusilla*
304 (Prasinophyceae) dominates the eukaryotic picoplankton in the western English Channel. *Appl. Environ.*
305 *Microbiol.* 70:4064–72.
- 306 Obayashi, Y. & Tanoue, E. 2002. Growth and mortality rates of phytoplankton in the northwestern North Pacific
307 estimated by the dilution method and HPLC pigment analysis. *J. Exp. Mar. Bio. Ecol.* 280:33–52.
- 308 R Development Core Team 2013. R: A Language and Environment for Statistical Computing.
- 309 Roesler, C.S., Culbertson, C.W., Etheridge, S.M., Goericke, R., Kiene, R.P., Miller, L.G. & Oremland, R.S. 2002.
310 Distribution, production, and ecophysiology of *Picocystis* strain ML in Mono Lake, California. *Limnol.*
311 *Oceanogr.* 47:440–52.
- 312 Sarada, R., Tripathi, U. & Ravishankar, G. 2002. Influence of stress on astaxanthin production in *Haematococcus*
313 *pluvialis* grown under different culture conditions. *Process Biochem.* 37:623–7.
- 314 Schlüter, L., Møhlenberg, F., Havskum, H. & Larsen, S. 2000. The use of phytoplankton pigments for identifying
315 and quantifying phytoplankton groups in coastal areas: Testing the influence of light and nutrients on
316 pigment/chlorophyll a ratios. *Mar. Ecol. Prog. Ser.* 192:49–63.
- 317 Shi, X.L., Marie, D., Jardillier, L., Scanlan, D.J. & Vaultot, D. 2009. Groups without cultured representatives
318 dominate eukaryotic picophytoplankton in the oligotrophic South East Pacific Ocean. *PLoS One.* 4.
- 319 Simon, N., Barlow, R.G., Marie, D., Partensky, F. & Vaultot, D. 1994. Characterization of oceanic photosynthetic
320 picoeukaryotes by flow cytometry analysis. *J. Phycol.* 30:922–35.
- 321 Subirana, L., Péquin, B., Michely, S., Escande, M.L., Meilland, J., Derelle, E., Marin, B. et al. 2013. Morphology,
322 genome plasticity, and phylogeny in the genus *Ostreococcus* reveal a cryptic species, *O. mediterraneus* sp.
323 nov. (Mamiellales, Mamiellophyceae). *Protist.* 164:643–59.

324 Throndsen, J. and Kristiansen, S. 1991. *Micromonas pusilla* (Prasinophyceae) as part of pico- and nanoplankton
325 communities of the Barents Sea. *Polar Res.* 10:201–7.

326 Viprey, M., Guillou, L., Ferréol, M. & Vaultot, D. 2008. Wide genetic diversity of picoplanktonic green algae
327 (Chloroplastida) in the Mediterranean Sea uncovered by a phylum-biased PCR approach. *Environ. Microbiol.*
328 10:1804–22.

329 Wang, B., Zarka, A., Trebst, A. & Boussiba, S. 2003. Astaxanthin accumulation in *Haematococcus pluvialis*
330 (Chlorophyceae) as an active photoprotective process under high irradiance. *J. Phycol.* 39:1116–24.

331 Wilhelm, C. & Lenarz-Weiler, I. 1987. Energy transfer and pigment composition in three chlorophyll b-containing
332 light-harvesting complexes isolated from *Mantoniella squamata* (Prasinophyceae), *Chlorella fusca*
333 (Chlorophyceae) and *Sinapis alba*. *Photosynth. Res.* 13:101–11.

334 Zapata, M., Rodriguez, F. & Garrido, J.L. 2000. Separation of chlorophylls and carotenoids from marine
335 phytoplankton: a new HPLC method using a reversed phase C-8 column and pyridine-containing mobile
336 phases. *Mar. Ecol. - Prog. Ser.* 195:29–45.

337

338

339

List of tables

Table 1. Characteristics of the strains used in this study. RCC refers to the Roscoff Culture Collection (<http://www.roscoff-culture-collection.org/>).

Table 2. Concentration of Chl *a* per cell, ratios of pigment to Chl *a* concentration and contribution to total carotenoids (in italics) for thirteen strains of prasinophytes clade VII (Chlorophyta) grown at $140 \mu\text{E.m}^{-2}.\text{s}^{-1}$.

List of supplementary tables

Table S1. Chromatographic retention times and spectral characteristics of the major pigments for strains of prasinophytes clade VII.

Table S2. Ratios of pigment to Chl *a* concentration and contribution to total carotenoids (in italics) for nine strains of prasinophytes clade VII under three light intensities.

List of Figures

Figure 1. Cumulative ratios of Chl *b* and of four major carotenoids (lutein *a*, violaxanthin, zeaxanthin, astaxanthin) to Chl *a* for 13 strains of prasinophytes clades VII at 140 $\mu\text{E.m}^{-2}.\text{s}^{-1}$. Strains are ordered by sub-clades (A, B, C) and depth of isolation (surface, deep chlorophyll maximum-DCM). RCC287a correspond to the composition reported by Latasa *et al.* (2004) for this strain.

Figure 2. Change in pigment to Chl *a* ratios for Chl *b* and five major carotenoids in nine strains of prasinophytes clade VII under three light intensities. Top: pigments involved in light capture. Bottom: pigments involved in photoprotection. Solid lines correspond to sub-clade VIIA and dashed lines to VIIB. Open symbols correspond to surface strains, closed ones to DCM strains and grey to unknown depth of isolation.

Figure 3. Principal component analysis using the pigment to Chl *a* ratios as variables for the strains grown at 3 light levels (Table S2). Top. Variables. Bottom. Samples. Circles correspond to clade VIIA and triangles to clade VIIB. Closed symbols correspond to low light, grey symbols to medium light and open symbols to high light.

List of Supplementary Figures

Figure S1. Change in pigment to Chl *a* ratios for Chl *b* and four minor carotenoids in nine strains of prasinophytes clade VII under three light intensities. Solid lines correspond to clade VIIA and dashed lines to VIIB. Open symbols correspond to surface strains, closed ones to DCM strains and grey to unknown depth of isolation.

Table 1

RCC	Sub-clade	Strain name	Other names	Ocean origin	Region origin	Depth Isolation (m)
15	A	CCMP 1205		NA		NA
287	A	NOUM15	NOUM97015	Pacific Ocean	Equatorial Pacific	120
719	A	IndianOcean_45-8		Indian Ocean		76
856	A	Biosope_42 A2	CCMP3325	Pacific Ocean	Marquesas islands	10
857	A	Biosope_40 A2		Pacific Ocean	Marquesas islands	10
996	A	Biosope_46 B4S		Pacific Ocean	South East Pacific	100
998	A	Biosope_46 C3S	NIES2676, CCMP3334	Pacific Ocean	South East Pacific	100
1124	A	PAP_AD	PAP_Ludwig_AI	Atlantic Ocean	North Atlantic, PAP site	10
3374	A	CCMP 2152	A7831	Pacific Ocean	Hawaii	NA
3376	A	CCMP 2113	A9533	Pacific Ocean		85
2337	B	JST MH335	MH335, NIES2756	Pacific Ocean	Iki Island	0
2339	B	JST MH340	MH340, NIES2758, CCMP3360	Pacific Ocean	Iki Island	0
3402	C	CCMP 1897	SFBB	Pacific Ocean	San Francisco Bay	0

Remark : Ordered by clade and then RCC number

Table 2

Strain	sub - clade	Light μE.m ⁻² .s ⁻¹	fg Chl <i>a</i> /cell	Chl <i>b</i> /Chl <i>a</i>	Chlide <i>a</i> /Chl <i>a</i>	Chlide <i>b</i> /Chl <i>a</i>	Sum of carotenoids /Chl <i>a</i>	Carotenoids																	
								Loroxanthin		Neoxanthin		Violaxanthin		Astaxanthin		Antheraxanthin		Zeaxanthin		Lutein <i>a</i>		ββ - carotene		βε - carotene	
								/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%
RCC15	A	140	20.3	0.90	0.066	0.018	1.32	0.058	4.4	0.003	0.2	0.327	24.8	0.165	12.5	0.043	3.3	0.103	7.8	0.376	28.4	0.154	11.7	0.093	7.0
RCC287	A	140	5.0	0.99	0.106	0	1.53	0.024	1.6	0.167	10.9	0.572	37.3	0.210	13.7	0.024	1.6	0.043	2.8	0.363	23.7	0.051	3.3	0.079	5.2
RCC719	A	140	14.9	0.68	0.000	0	1.93	0.035	1.8	0.086	4.4	0.617	32.0	0.264	13.7	0.000	0.0	0.478	24.8	0.193	10.0	0.107	5.5	0.149	7.7
RCC856	A	140	23.4	0.86	0.000	0	2.08	0.113	5.4	0.079	3.8	0.291	14.0	0.570	27.4	0.042	2.0	0.310	14.9	0.461	22.2	0.075	3.6	0.136	6.5
RCC857	A	140	4.1	1.00	0.000	0	1.40	0.026	1.9	0.122	8.7	0.534	38.2	0.187	13.4	0.024	1.7	0.094	6.7	0.291	20.8	0.056	4.0	0.064	4.5
RCC996	A	140	51.6	0.93	0.074	0	1.26	0.043	3.4	0.095	7.5	0.119	9.5	0.191	15.2	0.039	3.1	0.245	19.5	0.364	28.9	0.091	7.2	0.074	5.9
RCC998	A	140	26.1	0.78	0.000	0	1.90	0.014	0.8	0.117	6.1	0.877	46.1	0.157	8.2	0.058	3.0	0.168	8.8	0.285	15.0	0.095	5.0	0.131	6.9
RCC1124	A	140	8.6	0.96	0.099	0	1.47	0.000	0.0	0.116	7.9	0.525	35.7	0.148	10.1	0.020	1.4	0.086	5.8	0.399	27.1	0.095	6.5	0.082	5.6
RCC3374	A	140	4.1	0.73	0.229	0	1.84	0.012	0.7	0.082	4.5	0.272	14.8	0.778	42.3	0.041	2.3	0.118	6.4	0.332	18.1	0.059	3.2	0.146	7.9
RCC3376	A	140	4.1	0.81	0.105	0	1.53	0.008	0.5	0.112	7.3	0.572	37.5	0.324	21.3	0.054	3.6	0.129	8.5	0.048	3.1	0.085	5.6	0.194	12.7
RCC2337	B	140	4.4	0.88	0.394	0.236	2.19	0.031	1.4	0.137	6.3	0.504	23.1	0.302	13.8	0.051	2.3	0.154	7.0	0.706	32.3	0.061	2.8	0.240	11.0
RCC2339	B	140	14.6	0.62	0.000	0	1.52	0.026	1.7	0.078	5.1	0.613	40.4	0.045	3.0	0.024	1.6	0.074	4.8	0.302	19.9	0.028	1.8	0.330	21.7
RCC3402	C	140	131.1	0.26	0.016	0	0.54	0.000	0.0	0.048	8.9	0.014	2.7	0.000	0.0	0.000	0.0	0.071	13.1	0.200	37.0	0.016	3.0	0.191	35.4
RCC287 ^a	A	100	Nd	1.31	Nd	Nl	0.76	Nd	Nd	0.074	9.7	0.131	17.3	Nd	Nd	0.051	6.7	0.042	5.5	0.382	50.3	0.026	3.4	0.053	7.0

nd: Not determined

^aValues reported by Latasa *et al.* 2004

Table S1

Pigment	Abbreviation	Zapata et al. (2000)		Garrido et al. (2009)	
		Rt (min)	Vis maxima (nm)	Rt (min)	Vis maxima (nm)
Chlorophyllide <i>b</i>	Chlide <i>b</i>	6.21	466, 647		
Chlorophyllide <i>a</i>	Chlide <i>a</i>	11.43	431, 665		
Loroxanthin	Lor	20.18	443, 468	4.81	448, 475
Neoxanthin	Neo	20.18	443, 468	5.07	438, 467
Violaxanthin	Viola	21.92	443, 472	5.39	442, 470
Astaxanthin	Asta	22.94	481	5.63	478
Antheraxanthin	Anth	25.76	447, 476		
Zeaxanthin	Zea	27.85	454, 481		
Lutein	Lut	28.03	447, 476		
Chlorophyll <i>b</i>	Chl <i>b</i>	32.04	463, 649		
Chlorophyll <i>a</i>	Chl <i>a</i>	33.6	432, 664		
$\beta\epsilon$ -carotene	$\beta\epsilon$ -car	35.82	449, 476		
$\beta\beta$ -carotene	$\beta\beta$ -car	36.23	454, 479		

Table S2

			Carotenoids																						
Strain	sub - clade	Light μE/m2/s	FCM																						
			Mean FL3-A /cell	fg Chl a / cell	Chl b	Chlide a	Chlide b	Lor	Lor %	Neo	Neo%	Viola	Viol%	Asta	Asta%	Anth	Anth%	Zea	Zea%	Lut	Lut%	ββ - car	ββ - car%	βε - car	βε - car%
RCC287	A	140	97 734	4.99	0.986	0.106	0.000	0.024	1.6	0.167	10.9	0.572	37.3	0.210	13.7	0.024	1.6	0.043	2.8	0.363	23.7	0.051	3.3	0.079	5.2
		65	134 088	36.28	1.148	0.077	0.000	0.044	2.9	0.138	9.1	0.512	33.9	0.243	16.1	0.024	1.6	0.059	3.9	0.389	25.8	0.062	4.1	0.039	2.6
		14	187 046	20.13	1.041	0.000	0.000	0.071	7.4	0.100	10.5	0.282	29.5	0.085	8.9	0.019	2.0	0.032	3.3	0.230	24.1	0.089	9.3	0.047	4.9
RCC719	A	140	90 702	14.89	0.683	0.000	0.000	0.035	1.8	0.086	4.4	0.617	32.0	0.264	13.7	0.000	0.0	0.478	24.8	0.193	10.0	0.107	5.5	0.149	7.7
		65	101 762	22.74	0.832	0.000	0.000	0.066	3.9	0.118	6.9	0.722	42.7	0.137	8.1	0.044	2.6	0.194	11.5	0.204	12.1	0.140	8.3	0.069	4.1
		14	214 898	26.24	0.766	0.000	0.000	0.096	7.6	0.084	6.6	0.663	52.1	0.069	5.4	0.020	1.6	0.091	7.1	0.061	4.8	0.139	10.9	0.049	3.9
RCC856	A	140	145 875	23.37	0.860	0.000	0.000	0.113	5.4	0.079	3.8	0.291	14.0	0.570	27.4	0.042	2.0	0.310	14.9	0.461	22.2	0.075	3.6	0.136	6.5
		65	83 423	58.62	0.882	0.000	0.000	0.018	0.9	0.111	5.6	0.461	23.2	0.459	23.1	0.049	2.5	0.248	12.5	0.438	22.1	0.086	4.3	0.117	5.9
		14	231 009	56.88	0.942	0.000	0.000	0.096	8.2	0.102	8.8	0.310	26.6	0.138	11.8	0.023	1.9	0.093	8.0	0.222	19.0	0.133	11.4	0.049	4.2
RCC857	A	140	104 853	4.10	1.000	0.000	0.000	0.026	1.9	0.122	8.7	0.534	38.2	0.187	13.4	0.024	1.7	0.094	6.7	0.291	20.8	0.056	4.0	0.064	4.5
		65	150 099	11.46	1.128	0.000	0.000	0.072	5.7	0.131	10.4	0.400	31.9	0.141	11.3	0.025	2.0	0.072	5.7	0.314	25.1	0.065	5.2	0.034	2.7
		14	179 776	11.90	1.120	0.000	0.000	0.145	13.6	0.130	12.2	0.287	27.0	0.070	6.6	0.029	2.7	0.082	7.7	0.191	18.0	0.092	8.7	0.036	3.4
RCC996	A	140	253 221	51.57	0.931	0.074	0.000	0.043	3.4	0.095	7.5	0.119	9.5	0.191	15.2	0.039	3.1	0.245	19.5	0.364	28.9	0.091	7.2	0.074	5.9
		65	321 893	120.69	1.028	0.000	0.000	0.116	9.8	0.106	9.0	0.056	4.7	0.163	13.8	0.026	2.2	0.224	19.0	0.315	26.7	0.117	9.9	0.057	4.8
		14	418 516	61.80	0.968	0.000	0.000	0.195	17.8	0.120	11.0	0.399	36.5	0.049	4.4	0.016	1.5	0.058	5.3	0.077	7.1	0.179	16.4	0.000	0.0
RCC3374	A	140	53 612	4.14	0.726	0.229	0.000	0.012	0.7	0.082	4.5	0.272	14.8	0.778	42.3	0.041	2.3	0.118	6.4	0.332	18.1	0.059	3.2	0.146	7.9
		65	105 492	9.87	0.890	0.110	0.000	0.063	4.8	0.094	7.1	0.192	14.5	0.405	30.5	0.038	2.9	0.085	6.4	0.337	25.4	0.058	4.3	0.055	4.1
		14	136 153	11.33	0.777	0.000	0.000	0.113	12.1	0.098	10.5	0.176	18.8	0.187	20.1	0.020	2.1	0.035	3.8	0.159	17.0	0.082	8.8	0.063	6.8
RCC3376	A	140	50 333	4.08	0.813	0.105	0.000	0.008	0.4	0.112	5.7	0.572	29.1	0.324	16.5	0.054	2.7	0.129	6.6	0.486	24.7	0.085	4.3	0.194	9.9
		65	74 109	2.70	0.830	0.142	0.000	0.006	0.3	0.126	6.7	0.447	23.9	0.261	14.0	0.073	3.9	0.177	9.5	0.505	27.1	0.093	5.0	0.181	9.7
		14	101 238	10.08	0.783	0.053	0.000	0.005	0.5	0.095	9.5	0.108	10.8	0.091	9.1	0.029	2.9	0.130	13.0	0.365	36.6	0.055	5.5	0.122	12.2
RCC2337	B	140	84 624	4.37	0.882	0.394	0.236	0.031	1.4	0.137	6.3	0.504	23.1	0.302	13.8	0.051	2.3	0.154	7.0	0.706	32.3	0.061	2.8	0.240	11.0
		65	76 671	64.32	1.043	0.250	0.172	0.064	4.2	0.129	8.4	0.233	15.3	0.245	16.1	0.020	1.3	0.072	4.7	0.560	36.7	0.067	4.4	0.136	8.9
		14	213 378	18.09	1.040	0.158	0.118	0.109	9.1	0.137	11.4	0.255	21.3	0.058	4.8	0.028	2.4	0.067	5.6	0.332	27.7	0.118	9.8	0.095	7.9
RCC2339	B	140	123 995	14.58	0.624	0.000	0.000	0.026	1.7	0.078	5.1	0.613	40.4	0.045	3.0	0.024	1.6	0.074	4.8	0.302	19.9	0.028	1.8	0.330	21.7
		65	178 752	37.66	0.726	0.040	0.000	0.051	3.6	0.100	7.1	0.524	37.4	0.000	0.0	0.032	2.3	0.102	7.3	0.287	20.5	0.004	0.3	0.303	21.6
		14	223 724	75.86	0.736	0.000	0.000	0.051	5.8	0.076	8.6	0.264	29.8	0.031	3.5	0.025	2.9	0.051	5.8	0.193	21.8	0.033	3.7	0.160	18.1




Figure 1




Figure 2





Figure 3




Figure S1.